



Cambridge IGCSE™

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

BIOLOGY

0610/52

Paper 5 Practical Test

October/November 2025

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].

For Examiner's Use

1	
2	
3	
Total	

This document has **16** pages. Any blank pages are indicated.

- 1 You are going to investigate the rate of respiration in yeast using different concentrations of glucose.

You will use a 5% glucose solution to make a 1% glucose solution and a 4% glucose solution.

Read all the instructions but DO NOT DO THEM until you have drawn a table for your results in the space provided in 1(a)(i).

You should use the safety equipment provided while you are doing the practical work.

Step 1 Label two small test-tubes **1%** and **4%**. Put the small test-tubes in the test-tube rack.

Step 2 Put 1 cm³ of 5% glucose solution into the small test-tube labelled **1%**.

Put 4 cm³ of 5% glucose solution into the small test-tube labelled **4%**.

Step 3 Put 4 cm³ of distilled water into the small test-tube labelled **1%**.

Put 1 cm³ of distilled water into the small test-tube labelled **4%**.

Step 4 Raise your hand when you are ready for hot water to be added to the water-bath.

Step 5 Pour yeast suspension into the small test-tube labelled **1%** until it is completely full.

Step 6 Keep the small test-tube upright and carefully place a large test-tube over the top of the small test-tube labelled **1%**. Gently push the small test-tube up to the top of the large test-tube, as shown in Fig. 1.1A.

Step 7 Keep the small test-tube in place with your finger and then quickly turn the test-tubes so they are inverted, as shown in Fig. 1.1B. It does not matter if a small amount of air remains at the top of the small test-tube when it is inverted.




Fig. 1.1



- Step 8 Put the large test-tube and the small test-tube labelled **1%** into the water-bath, as shown in Fig. 1.1C.
- Step 9 Start the stop-clock and wait for five minutes.
- Step 10 After five minutes, stop the stop-clock and use the ruler to measure the height of the yeast suspension in the **large** test-tube, as shown in Fig. 1.2. Record the measurement in your table in **1(a)(i)**.




Fig. 1.2

- Step 11 Remove the test-tubes from the water-bath, and place them back in the test-tube rack.
- Step 12 Pour yeast suspension into the small test-tube labelled **4%** until it is completely full.
- Step 13 Repeat steps 6 to 10 with the small test-tube labelled **4%** and the remaining empty large test-tube.





(a) (i) Prepare a table and record your results.

[4]

(ii) State a conclusion for your results.

.....

.....

..... [1]

(iii) State the independent and dependent variables in this investigation.

independent variable

.....

dependent variable

..... [2]

(iv) Describe how the temperature of the water-bath could be maintained.

.....

.....

..... [1]



- (v) Identify **one** possible source of error when measuring the height of the yeast suspension in step 10.

.....

 [1]

- (b) A student extended the investigation by using more concentrations of glucose solution.

Calculate the volumes of 5% glucose solution and distilled water that are needed to make 5 cm³ of a 3% glucose solution.

volume of 5% glucose solution cm³

volume of distilled water cm³
 [1]

- (c) Sports drinks often contain glucose, which is a reducing sugar.

State how you could test a sample of a sports drink for reducing sugar.

.....

 [2]

- (d) (i) The ethanol emulsion test can be used to show that a sample of food contains fat.

State the result of a positive test.

..... [1]

- (ii) Suggest **one** safety precaution that should be followed when using ethanol.

.....

 [1]

[Total: 14]





[6]



- 3 (a) Fig. 3.1 is a photograph of a necklace starfish, *Fromia monilis*.




Fig. 3.1





- (i) Draw a large diagram of the necklace starfish shown in Fig. 3.1.

[4]

- (ii) Line **PQ** in Fig. 3.1 represents the diameter of the necklace starfish.

Measure the length of line **PQ** in Fig. 3.1.

length of **PQ** mm

Calculate the actual diameter of the necklace starfish using the formula and your measurement.

$$\text{magnification} = \frac{\text{length of line PQ in Fig. 3.1}}{\text{actual diameter of the necklace starfish}}$$

Give your answer to **two** significant figures.

Space for working.

..... mm
[3]




- (iii) Fig. 3.2 is a photograph of a spiny starfish, *Marthasterias glacialis*. Fig. 3.3 is a photograph of a necklace starfish.

The magnifications of the photographs are **not** the same.



spiny starfish

Fig. 3.2



necklace starfish

Fig. 3.3

State **two** ways the spiny starfish in Fig. 3.2 differs from the necklace starfish in Fig. 3.3, and **one** way it is similar.

Do **not** include references to size in your answer.

difference 1

.....

difference 2

.....

similarity 1

.....

[3]



(b) Seven-armed starfish live in the sea.

Scientists investigated the distribution of seven-armed starfish, *Luidia ciliaris*, in one region.

- Photographs of 20 m² areas of seabed were taken at different depths.
- The depth of the water in each area was measured when the water was at its highest level (high tide).
- The numbers of seven-armed starfish in each photograph were counted.
- All photographs were taken on the same day in July.

- (i) State **two** ways that the method used by the scientists ensured that the numbers of seven-armed starfish counted at different depths could be compared.

1

.....

2

.....

[2]

The results of the investigation are shown in Table 3.1.






Table 3.1

depth/m	number of seven-armed starfish counted
0.0–9.9	337
10.0–19.9	1486
20.0–29.9	1243
30.0–39.9	241
40.0–49.9	121

(ii) Plot a histogram on the grid of the data in Table 3.1.



- (iii) State the depth range in Table 3.1 at which most seven-armed starfish were found.

.....m [1]

- (iv) Using the information in Table 3.1, calculate the total number of seven-armed starfish that were photographed at all depths, and use this value to calculate the percentage of starfish that were photographed in the 0.0–9.9m range.

Express the percentage as a whole number.

Space for working.

total number of seven-armed starfish photographed

percentage of seven-armed starfish photographed at 0.0–9.9m%
[3]

[Total: 20]









Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

