



Mark Scheme (Results)

January 2020

Pearson Edexcel International GCSE in
Chemistry (4CH1)
Paper 1C

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2020

Publications Code 4CH1_1C_msc_2020305

All the material in this publication is copyright

© Pearson Education Ltd 2020


General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	(i) argon / helium	ACCEPT Ar / He	1
	(ii) nitrogen	ACCEPT N ₂	1
	(iii) carbon dioxide	ACCEPT CO ₂	1
	(iv) carbon dioxide	ACCEPT CO ₂	1
(b)	(i) $S + O_2 \rightarrow SO_2$		1
	(ii) acid rain	ACCEPT an adverse effect of acid rain e.g. kills fish, damages plants, corrodes limestone/marble buildings/statues etc. IGNORE toxic/pollutant	1
			Total 6

Question number	Answer	Notes	Marks										
2 (a)	<table><tr><td>name of the part of the atom labelled Z</td><td>nucleus</td></tr><tr><td>number of protons in this atom</td><td>12</td></tr><tr><td>number of the group that contains this element</td><td>2</td></tr><tr><td>number of the period that contains this element</td><td>3</td></tr><tr><td>charge on the ion formed from this atom</td><td>2+</td></tr></table>	name of the part of the atom labelled Z	nucleus	number of protons in this atom	12	number of the group that contains this element	2	number of the period that contains this element	3	charge on the ion formed from this atom	2+	ACCEPT +2 / Mg ²⁺	5
name of the part of the atom labelled Z	nucleus												
number of protons in this atom	12												
number of the group that contains this element	2												
number of the period that contains this element	3												
charge on the ion formed from this atom	2+												
(b)	- calculate sum of mass numbers multiplied by percentage abundances - divide answer by 100 - give answer to one decimal place Example calculation M1 (24 x 79.2) + (25 x 10.0) + (26 x 10.8) OR 2431.6 M2 2431.6 ÷ 100 OR 24.316 M3 24.3	REJECT if correct working given but incorrectly evaluated ALLOW ECF from M1 (24 x 0.792) + (25 x 0.100) + (26 x 0.108) OR 24.316 with or without working scores M1 and M2 ALLOW ECF from M2 if calculated answer is to 1dp	3										
			Total 8										

Question number	Answer	Notes	Marks
3 (a)	galvanising	ACCEPT galvanisation	1
(b) (i)	rust		1
(ii)	M1 oxygen / air M2 water	ACCEPT O ₂ IGNORE O ACCEPT H ₂ O/moisture ACCEPT in either order	2
(c) (i)	(a reaction which) gives out / produces / releases heat (energy) / thermal energy	IGNORE energy without mention of heat or thermal	1
(ii)	An explanation that links the following two points M1 aluminium/Al is more reactive than iron/Fe M2 (because) aluminium/Al displaces iron/Fe (from its oxide)	ACCEPT aluminium/Al is higher in reactivity series than iron/Fe ACCEPT reverse argument ALLOW replaces/takes place of	2
(iii)	An explanation that links the following three points M1 aluminium is oxidised and iron/iron oxide is reduced M2 aluminium gains oxygen M3 iron oxide/iron loses oxygen	ALLOW both oxidation and reduction occur ALLOW aluminium/Al loses electrons ALLOW iron <u>ions</u> /Fe ³⁺ gains electrons ALLOW correct references to changes in oxidation number for M2 and M3	3
			Total 10

Question number	Answer	Notes	Marks																
4 (a) (i)	<table border="1"> <tr> <td></td><td>Mg^{2+}</td><td>Al^{3+}</td><td>NH_4^+</td></tr> <tr> <td>S^{2-}</td><td>MgS</td><td>Al_2S_3</td><td>$(\text{NH}_4)_2\text{S}$</td></tr> <tr> <td>NO_3^-</td><td>$\text{Mg}(\text{NO}_3)_2$</td><td>$\text{Al}(\text{NO}_3)_3$</td><td>NH_4NO_3</td></tr> <tr> <td>CO_3^{2-}</td><td>MgCO_3</td><td>$\text{Al}_2(\text{CO}_3)_3$</td><td>$(\text{NH}_4)_2\text{CO}_3$</td></tr> </table>		Mg^{2+}	Al^{3+}	NH_4^+	S^{2-}	MgS	Al_2S_3	$(\text{NH}_4)_2\text{S}$	NO_3^-	$\text{Mg}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$	NH_4NO_3	CO_3^{2-}	MgCO_3	$\text{Al}_2(\text{CO}_3)_3$	$(\text{NH}_4)_2\text{CO}_3$	1 mark for each correct formula	3
	Mg^{2+}	Al^{3+}	NH_4^+																
S^{2-}	MgS	Al_2S_3	$(\text{NH}_4)_2\text{S}$																
NO_3^-	$\text{Mg}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$	NH_4NO_3																
CO_3^{2-}	MgCO_3	$\text{Al}_2(\text{CO}_3)_3$	$(\text{NH}_4)_2\text{CO}_3$																
(ii)	ammonium nitrate		1																
(b) (i)	M1 electrostatic (force of) attraction M2 between oppositely charged ions	ALLOW electrostatic force ACCEPT between positive and negative ions ACCEPT between cations and anions	2																
(ii)	 M1 correct electron arrangement of both sodium ions M2 correct electron arrangement of the oxide ion M3 correct charges on all ions (with or without brackets)	If only outer shells shown correctly scores 1 mark ACCEPT dots in place of crosses or any combination of dots and crosses for M1 and M2	3																
Total 9																			

Question number	Answer	Notes	Marks
5 (a) (i)	S		1
(ii)	T and U		1
(iii)	U		1
(b)	A description that makes reference to the following three points M1 (add) bromine water M2 no change / stays orange M3 (bromine water) decolourised / changes (from orange) to colourless	ACCEPT Br₂ (aq) ALLOW no reaction If initial colour of bromine water is given in M2 or M3 it must be correct -ALLOW any combination of orange/yellow/brown - but penalise once only If bromine given for M1 then in M2 and M3 allow any combination of red/orange/brown/yellow M2 and M3 dep on bromine water/bromine in M1 If no reagent and correct M2 and M3 - score 1 if incorrect reagent and correct M2 and M3 score 0 IGNORE clear REJECT discoloured ALLOW M1 acidified potassium manganate(VII) M2 no change/stays purple M3 decolourised / goes colourless	3

Question number	Answer	Notes	Marks
5 (c)	Any two of the following points M1 (can be represented by a) general formula M2 each member differs from the next by a CH₂ group OWTTE M3 (each member has) same functional group M4 (each member has) similar/same chemical properties / similar/same (chemical) reactions M5 trend in physical properties (between successive members)	ACCEPT react in similar/same way ACCEPT named physical property, e.g. boiling point REJECT similar/same physical properties	2
(d) (i)	but-1-ene	ALLOW 1-butene	1
(ii)	Either $ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & & \text{H} \\ & & & & & & & \\ \text{H} & - \text{C} & - & \text{C} & = & \text{C} & - & \text{C} & - \text{H} \\ & & & & & & & \\ & \text{H} & & & & & & \text{H} \end{array} $ Or $ \begin{array}{ccccc} & \text{H} & & & \text{H} \\ & & & & / \\ \text{H} & - \text{C} & - & \text{C} & = & \text{C} \\ & & & & & \backslash \\ & \text{H} & & & & \text{H} \\ & & & & & / \\ & \text{H} & - & \text{C} & - & \text{H} \\ & & & & & \\ & & & \text{H} & & \end{array} $	ACCEPT cis or trans isomer REJECT displayed formulae of cyclic alkanes	1

Question number	Answer	Notes	Marks																
5 (e) (i)	- Divide percentages by relative atomic masses - Divide results by smallest value to obtain ratio Example calculation <table><tr><td>M1</td><td>C</td><td>H</td><td>F</td></tr><tr><td></td><td>$\frac{36.36}{12}$</td><td>$\frac{6.06}{1}$</td><td>$\frac{57.58}{19}$</td></tr><tr><td>M2</td><td>$\frac{3.03}{3.03}$</td><td>$\frac{6.06}{3.03}$</td><td>$\frac{3.03}{3.03}$</td></tr><tr><td>OR</td><td>1</td><td>2</td><td>1</td></tr></table>	M1	C	H	F		$\frac{36.36}{12}$	$\frac{6.06}{1}$	$\frac{57.58}{19}$	M2	$\frac{3.03}{3.03}$	$\frac{6.06}{3.03}$	$\frac{3.03}{3.03}$	OR	1	2	1	0 marks if division by atomic numbers or upside down calculation	2
	M1	C	H	F															
	$\frac{36.36}{12}$	$\frac{6.06}{1}$	$\frac{57.58}{19}$																
M2	$\frac{3.03}{3.03}$	$\frac{6.06}{3.03}$	$\frac{3.03}{3.03}$																
OR	1	2	1																
(ii)	- divide relative molecular mass by empirical formula mass - correct molecular formula Example calculation <table><tr><td>M1</td><td>$\frac{66}{12 + 2 + 19}$</td><td>OR</td><td>$\frac{66}{33}$</td><td>OR</td><td>2</td></tr><tr><td>M2</td><td>C₂H₄F₂</td><td></td><td></td><td></td><td></td></tr></table>	M1	$\frac{66}{12 + 2 + 19}$	OR	$\frac{66}{33}$	OR	2	M2	C ₂ H ₄ F ₂					ACCEPT symbols in any order correct answer without working scores 2 marks. 2CH₂F scores 1	2				
M1	$\frac{66}{12 + 2 + 19}$	OR	$\frac{66}{33}$	OR	2														
M2	C ₂ H ₄ F ₂																		
Total 14																			

Question number	Answer	Notes	Marks							
6	(a)	zinc + hydrochloric acid → zinc chloride + hydrogen	ACCEPT fully correct chemical equation	1						
	(b)	<table><tr><td>temperature in °C after adding zinc</td><td>22.4</td></tr><tr><td>temperature in °C before adding zinc</td><td>17.7</td></tr><tr><td>temperature change in °C</td><td>4.7</td></tr></table>	temperature in °C after adding zinc	22.4	temperature in °C before adding zinc	17.7	temperature change in °C	4.7	If readings are correct but in wrong order award 1 mark for M1 and M2	3
		temperature in °C after adding zinc	22.4							
		temperature in °C before adding zinc	17.7							
	temperature change in °C	4.7								
	M1 22.4									
	M2 17.7									
		M3 (+)4.7	ALLOW ECF for M3 if M1 and/or M2 incorrect If answers not given to nearest 0.1 °C penalise once only							
	(c)	(i)	An explanation that links any two of the following points		2					
			M1 polystyrene is an insulator	ALLOW is not a (good) conductor of heat ALLOW is a poor conductor of heat						
M2 (so) reduces heat loss			ALLOW prevents heat loss ALLOW keeps heat in							
		M3 temperature rise/change/reading will be closer to true value OWTTE	ALLOW temperature rise/change/reading will be more accurate/valid							
(ii)		Any three from								
		M1 amount/mass of metal	ALLOW size / surface area of metal							
		M2 concentration of acid								
		M3 volume of acid	ALLOW amount of acid							
		M4 (speed/time of) stirring								
		M5 external / room temperature	ALLOW initial /starting temperature	3						

Question number	Answer	Notes	Marks
6 (d)	(i) no reaction (occurred between copper and hydrochloric acid)	IGNORE copper is unreactive ALLOW copper is less reactive than hydrogen	1
	(ii) Any value between 1.5 and 5.0 °C inclusive		1
	(iii) most reactive magnesium zinc iron tin copper	ACCEPT symbols	1
			Total 12

Question number	Answer	Marks
7 (a)	(i) B bromine	1
	(ii) A is incorrect as astatine is a solid C is incorrect as chlorine is a gas D is incorrect as iodine is a solid	1
	(iii) C chlorine (as it is pale green) A is incorrect as astatine is black B is incorrect as bromine is brown C is incorrect as iodine is dark grey A astatine B is incorrect as bromine is more reactive than astatine C is incorrect as chlorine is more reactive than astatine D is incorrect as iodine is more reactive than astatine	1

(b) (i)	M1 (colourless solution turns) brown M2 (solution stays) brown / no change	ALLOW no reaction	2
(ii)	bromine would not react with (sodium) bromide / bromine cannot displace itself OWTTE	ALLOW bromine cannot react with itself ALLOW both contain bromine/same element/same halogen ALLOW because no reaction would occur REJECT bromine cannot displace bromide	1
(iii)	$\text{Br}_2 + 2\text{NaI} \rightarrow 2\text{NaBr} + \text{I}_2$	ACCEPT correct ionic equation $\text{Br}_2 + 2\text{I}^- \rightarrow 2\text{Br}^- + \text{I}_2$ ALLOW multiples and fractions	1

Question number	Answer	Notes	Marks
7 (c)	A description that makes reference to the following 6 points Test for cation M1 add sodium hydroxide (solution) M2 if blue precipitate forms solution contains copper(II) ion(s) / contains Cu^{2+} / is a copper compound M3 if green precipitate forms solution contains iron(II) ion(s) / contains Fe^{2+} / is an iron compound	ALLOW ammonia solution IGNORE qualifiers REJECT other colours IGNORE qualifiers REJECT other colours If no reagent or incorrect reagent but correct M2 and M3 score 1 ALLOW M1 flame test or description of flame test M2 if blue-green (flame) solution contains copper(II) ion(s) / contains Cu^{2+} / is a copper compound No M3 for this test ALLOW M1 addition of suitable metal above Cu in reactivity series M2 brown/pink/pink-brown solid forms No M3 for this test	


	Test for anion M4 add silver nitrate (solution) M5 if white precipitate forms solution contains chloride ion(s) / contains Cl^- / is a chloride M6 if cream precipitate forms solution contains bromide ion(s) / contains Br^- / is a bromide	IGNORE addition of nitric acid REJECT addition of hydrochloric acid for M4 If no reagent or incorrect reagent but correct M5 and M6 score 1 ALLOW M4 add chlorine water (to solution) M5 if turns orange/yellow/brown solution contains bromide ion(s) / contains Br^- / is a bromide No M6 for this test	Total 13
--	--	--	------------------------

Question number	Answer	Notes	Marks
8 (a) (i)	sublimation / subliming		1
(ii)	M1 (add to/bubble into) limewater M2 (limewater) turns cloudy/milky	ACCEPT forms white precipitate M2 DEP M1	2
(b)	An explanation that links the following two points M1 weak forces (of attraction) between molecules / weak intermolecular forces (of attraction) M2 little energy needed to overcome the (intermolecular) forces	ALLOW weak intermolecular bonds ALLOW weak intermolecular attractions IGNORE less energy ALLOW little energy needed to separate the molecules M2 DEP M1 correct or missing	2

(c)	Any explanation that links any three of the following points for diamond M1 each (carbon) atom is (covalently) bonded to four other (carbon) atoms M2 in a (giant) tetrahedral lattice /network / structure M3 the (covalent) bonds are (very) strong M4 (therefore) diamond is (very) hard (and so good for cutting tools) Any explanation that links any three of the following points for graphite M5 each (carbon) atom is (covalently) bonded to three other (carbon) atoms M6 (the structure is) in layers M7 weak forces (between layers) M8 (the layers can) slide over each other/ rub off M9 this makes graphite soft (so it can make marks on paper)	ALLOW each carbon has four bonds ALLOW 3D/rigid in place of tetrahedral ALLOW reference to lot of energy needed to break the (covalent) bonds ALLOW there are lots of/many (covalent) bonds ALLOW diamond is (very) strong If mention of intermolecular forces in diamond MAX 2 for diamond If mention of ions in diamond only M4 can be scored ALLOW sheets ALLOW slippery If mention of intermolecular forces in graphite MAX 2 for graphite If mention of ions in graphite only M9 can be scored	6 Total 11
-----	---	---	---------------------------------

Question number	Answer	Notes	Marks
9 (a)	to minimise/prevent (mass loss by) evaporation of the (liquid) fuel OWTTE	ALLOW to find mass of fuel used/burned	1
(b) (i)	soot/carbon	REJECT copper oxide	1
(ii)	An explanation that links the following two points. M1 incomplete combustion (occurs) M2 (because) the air/oxygen supply is limited OWTTE	ALLOW mark for soot/carbon if not seen in (i), unless copper oxide is mentioned in (i) If copper oxide in (i) ALLOW 1 mark for (because) copper reacts with oxygen (in air)	2
(c) (i)	- substitution into $Q = mc\Delta T$ - calculation of heat energy in Joules - conversion to kJ Example calculation M1 $Q = 100 \times 4.2 \times 30$ M2 = 12600 (J) M3 = 12.6 kJ	12600 (J) with no working scores M1 and M2 M2 ECF M1 ALLOW approximately = 13 kJ 12.6 kJ with no working scores 3	3

(ii)	• calculate the amount, in moles, of methanol • divide Q by the amount in moles • give the answer with the correct sign Example calculation M1 $0.96 \div 32$ OR 0.03 M2 $12.6 \div 0.03$ OR 420 (kJ/mol) M3 $- 420$ (kJ/mol)	ACCEPT $13 \div 0.03$ OR $430/433$ for M2 AND $- 430 / - 433$ for M3	3
------	---	---	---

Question number	Answer	Notes	Marks
9 (d) (i)	M1 all points plotted correctly M2 line of best fit drawn with a ruler 	does not need to start at (0,0)	2
(ii)	M1 straight line extrapolated up to 6 carbon atoms M2 value of ΔH read from their graph	ALLOW extra point shown at 6 carbon atoms negative sign needed	2
(iii)	The greater the number of carbon atoms (per molecule) the greater (the magnitude/ value of) ΔH	ALLOW ΔH is (directly) proportional to the number of carbon atoms per molecule ALLOW The greater the number of carbon atoms (per molecule) the more exothermic the ΔH value	1
Total 15			

Question number	Answer	Notes	Marks
10 (a) (i)	$4\text{NH}_3 + 5\text{O}_2 \rightleftharpoons 4\text{NO} + 6\text{H}_2\text{O}$	ACCEPT multiples and fractions	1
(ii)	reversible (reaction)	ACCEPT reaction that goes both ways / both forwards and backwards reactions occur IGNORE references to equilibrium	1
(iii)	to increase the rate of the reaction / to speed up the reaction OWTTE	IGNORE references to lowering the activation energy	1
(b)	$2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$	ACCEPT multiples and fractions	1
(c) (i)	- calculate M_r of NO_2 and HNO_3 - calculate the amount, in moles, of NO_2 - calculate the amount, in moles, of HNO_3 - calculate the mass in tonnes of HNO_3 Example calculation M1 M_r of $\text{NO}_2 = 46$ M_r of $\text{HNO}_3 = 63$ M2 $n(\text{NO}_2) = 11.5 \times 10^6 \div 46$ OR 250 000 (mol) M3 $n(\text{HNO}_3) = \frac{2 \times 25\,0000}{3}$ OR 167 000 / 170 000 M4 $(167\,000 \times 63 \text{ g}) = 10.5 \text{ (tonnes)}$	ALLOW working in megamoles i.e. $11.5 \div 46$ OR 0.25 ALLOW ECF from incorrect M_r of NO_2 calculator answer 166666.66 ALLOW working in megamoles i.e. $\frac{2 \times 0.25}{3}$ OR 0.167 / 0.17 ALLOW ECF from M2 10.5 (tonnes) with no working scores 4 ACCEPT 10.7 (if 170 000 used) ALLOW ECF from M3 ALLOW ECF from incorrect M_r of HNO_3	4
(ii)	can be (re)used in stage 2 / to make more nitrogen dioxide (in stage 2) / can be used to make more nitric acid	IGNORE can be recycled/reused unless qualified	1

Question number	Answer	Notes	Marks
10 (d)	- calculate the amount, in moles, of copper(II) nitrate - calculate the theoretical yield, in moles, of copper(II) nitrate - calculate the percentage yield Example calculation M1 $n\text{Cu}(\text{NO}_3)_2 \text{ formed} = 15.3 \div 187.5$ OR 0.0816 M2 theoretical $n\text{Cu}(\text{NO}_3)_2 = 0.200 \div 2$ OR 0.100 M3 (% yield) = $\frac{(0.0816 \times 100)}{(0.100)} = 81.6$ (%) Alternative method - calculate the theoretical yield, in moles, of copper(II) nitrate - calculate the theoretical mass of copper nitrate that should be formed - calculate the percentage yield Example calculation M1 theoretical $n\text{Cu}(\text{NO}_3)_2 = 0.200 \div 2$ OR 0.100 M2 theoretical mass of copper nitrate = $0.1 \times 187.5 = 18.75$ M3 (% yield) = $\frac{15.3}{18.75} \times 100 = 81.6$ (%)	ALLOW 0.082 ACCEPT 82 (%) Mark M3 CSQ on M1 and M2 40.8 scores 2 ALLOW 18.8 ACCEPT 82 (%) Mark M3 CSQ on M1 and M2 40.8 scores 2 81.6(%) with no working scores 3 marks	3 Total 12

