
CHEMISTRY

9701/22

Paper 2 AS Level Structured Questions

March 2019

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the March 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **9** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:


Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:


Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED

Question	Answer	Marks				
1(a)	strong triple bond / strong $\text{N}\equiv\text{N}$ OR high activation energy / E_{a} OR non-polar	1				
1(b)(i)	$3\text{Mg} + \text{N}_2 \rightarrow \text{Mg}_3\text{N}_2$	1				
1(b)(ii)	solid disappears	1				
1(c)(i)	(it is used to make) fertilisers	1				
1(c)(ii)	M1 CaO displaces NH_3 (from its salt / NH_4^+) M2 CaO is a stronger base / more basic (than NH_3)	2				
1(d)(i)	<table><tr><td>NO</td><td>NO_2</td></tr><tr><td>(+)2 / (+)II</td><td>(+)4 / (+)IV</td></tr></table>	NO	NO_2	(+)2 / (+)II	(+)4 / (+)IV	1
NO	NO_2					
(+)2 / (+)II	(+)4 / (+)IV					
1(d)(ii)	M1 $\frac{1}{2}\text{N}_2 + \text{O}_2 \rightarrow \text{NO}_2$ M2 $\text{Mg}(\text{NO}_3)_2 \rightarrow \text{MgO} + 2\text{NO}_2 + \frac{1}{2}\text{O}_2$	2				
1(d)(iii)	M1 $+82 (= E_{\text{O}=\text{O}} - 2E_{\text{N}\rightarrow\text{O}}) = (+)496 - 2 \times E_{\text{N}\rightarrow\text{O}}$ M2 $E_{\text{N}\rightarrow\text{O}} = \frac{1}{2} \times (496 - 82) = \frac{1}{2} \times 414 = 207 \text{ (kJ mol}^{-1}\text{)}$	2				


Question	Answer	Marks
1(e)	 M1 curly arrow from C=C to Nδ^+ AND curly arrow from N—Cl to Clδ^- M2 intermediate AND curly arrow from lone pair on Cl$^-$ to C(+)	2

Question	Answer	Marks
2(a)(i)	M1 ① mass of a molecule OR ② (weighted) average / mean mass of the molecules OR ③ mass of one mole of molecules M2 ① / ② compared to $\frac{1}{12}$ (the mass) of an atom of carbon-12 OR on a scale in which a carbon-12 atom / isotope has a mass of (exactly) 12 (units) ③ relative / compared to $\frac{1}{12}$ (the mass) of 1 mole of carbon-12 OR on a scale in which 1 mole of carbon-12 (atoms / isotope) has a mass of (exactly) 12 g	2

Question	Answer	Marks
2(a)(ii)	M1 identification of the IMF between F₂ molecules and between HCl molecules HCl has (permanent) dipoles and / or induced dipoles F₂ has induced dipoles M2 comparison of strength of IMF's in F₂ and HCl Intermolecular forces in HCl are stronger than F₂	2
2(a)(iii)	strong (electrostatic) forces of attraction between (oppositely charged) ions	1
2(a)(iv)	CaCO₃(s) + 2HF(aq) → CaF₂(aq) + CO₂(g) + H₂O(l) M1 species and balancing M2 state symbols	2
2(b)(i)	1s ² 2s ² 2p ⁶ 3s ² 3p ⁶	1
2(b)(ii)	M1 purple gas / vapour disappears M2 iodine is not a strong enough oxidising agent ORA	2
2(b)(iii)	M1 silver nitrate / AgNO₃ M2 yellow	2
2(b)(iv)	(aqueous) ammonia / NH ₃ (aq) / ammonium hydroxide / NH ₄ OH(aq)	1
2(c)(i)	 M1 bonding pairs correct M2 rest of molecule, incl. lone pairs.	2
2(c)(ii)	F ₂ + H ₂ O → HF + HOF	1


Question	Answer	Marks
2(c)(iii)	M1 labelled reactants AND products lower on right M2 labelled enthalpy change with correct arrow	2
2(c)(iv)	$\delta^+ \quad \delta^- \quad \delta^+ \quad \delta^-$ $\text{H}-\text{F} \vdots \vdots \vdots \vdots \vdots \vdots \text{H}-\text{F}$ M1 H-bond labelled / shown as distinct from H—F bond M2 correct sequence of three correct dipoles M3 lone pair on F in line with H-bond	3
2(d)(i)	$pV = nRT \therefore n = \frac{pV}{RT} = \frac{101\,325 \times 0.001}{8.31 \times 273} = 0.0447 \text{ mol}$ $\therefore M_r = \frac{m}{n} = \frac{4.13}{0.0447} = 92.4 \text{ or } 92.5$ M1 Use of $T = 273 \text{ K}$, $V = 0.001 \text{ m}^3$ and $p = 101325 \text{ Pa}$ M2 correct use of $pV = nRT$ using values from M1 M3 correct calculation of M_r using $4.13 \div$ moles from M2	3
2(d)(ii)	ClF_3	1

PUBLISHED

Question	Answer				Marks																
3(a)	<table><tr><td></td><td>P</td><td>Q</td><td>R</td></tr><tr><td>Na(s)</td><td>effervescence</td><td>no reaction</td><td>no reaction</td></tr><tr><td>2,4-DNPH</td><td>no reaction</td><td>orange ppt</td><td>orange ppt</td></tr><tr><td>acidified K₂Cr₂O₇(aq)</td><td>no reaction</td><td>no reaction</td><td>(turns) green</td></tr></table>					P	Q	R	Na(s)	effervescence	no reaction	no reaction	2,4-DNPH	no reaction	orange ppt	orange ppt	acidified K ₂ Cr ₂ O ₇ (aq)	no reaction	no reaction	(turns) green	3
	P	Q	R																		
Na(s)	effervescence	no reaction	no reaction																		
2,4-DNPH	no reaction	orange ppt	orange ppt																		
acidified K ₂ Cr ₂ O ₇ (aq)	no reaction	no reaction	(turns) green																		
3(b)	C ₅ H ₁₀ O + 2[H] → C ₅ H ₁₂ O				1																
3(c)	M1 geometric(al) M2 				2																
3(d)	M1 compound P M2 (absorption at) 2250 cm⁻¹ AND C≡N (stretch) M3 (absorption at) 3100–3700 cm⁻¹ AND O—H (stretch)				3																

Question	Answer	Marks
4(a)	3-chloroprop-1-ene	1
4(b)(i)	ultra-violet (light) / sun(light)	1
4(b)(ii)	$\text{CH}_2=\text{CHCH}_3 + \text{Cl}^\bullet \rightarrow \text{CH}_2=\text{CHCH}_2^\bullet + \text{HCl}$ OR $\text{C}_3\text{H}_6 + \text{Cl}^\bullet \rightarrow \text{C}_3\text{H}_5^\bullet + \text{HCl}$	1

PUBLISHED

Question	Answer	Marks
4(b)(iii)	free-radical (substitution) reactions are uncontrolled OR further chlorination / substitution occurs	1
4(b)(iv)	SOCl_2 OR PCl_5 OR PCl_3 OR <u>c</u> (oncentrated) HCl	1
4(c)(i)	cold, dilute acidified KMnO_4 / potassium manganate(VII)	1
4(c)(ii)	M1 catalyst M2 ethanoic acid / $\text{CH}_3\text{CO}_2\text{H}$	2
4(c)(iii)	nucleophilic substitution / $\text{S}_{\text{N}}2$	1
4(c)(iv)	 M1 hydrolysed nitrile on straight-chain 4C backbone M2 3,4-diol	2
4(d)	M1 major product formed from more stable intermediate / carbocation OR (intermediate has) 2° carbocation which is (more) stable M2 (positive) inductive effect / (+)I of alkyl groups (on the intermediate)	2